

Colonoscopy Prep Instructions – Extended Prep – Clenpiq/Plenvu/Bisacod



YOUR BOWEL PREP IS VERY IMPORTANT- Please read immediately and review instructions carefully! Only follow these MDTEC instructions for your prep.

If you do not follow these instructions and you are not prepped properly for your procedure, a cancellation fee will apply.

Prior to your procedure:		2 Days Prior	1 Day Prior	Procedure Day
<p>Your prescription is sent to your pharmacy after your pre-op call or office visit has been completed. Please pick up your prescription within 72 hours.</p> <p>Purchase Bisacodyl from the pharmacy.</p> <p>You will need to arrange for a responsible adult to drive you home. You cannot drive until the day after your procedure.</p> <p>A taxi, ride share, or shuttle is not an approved means of transportation unless you have a family member or friend with you.</p> <p>If you take Phentermine – stop taking this 10 days prior to your procedure.</p> <p>If you take iron, dietary, herbal or fiber supplements, stop taking these 7 days prior to your procedure.</p> <p>If you take a blood thinner-</p> <p>Please contact the prescribing doctor to obtain a blood thinner hold order. The office can fax this order to us at 410-972-2698.</p> <p>If you see a specialist: you can request that your most recent visit summary, labs, and test results from that office be faxed to us at 410-972-2698</p> <p>If you develop any covid, flu, or cold like symptoms within 2 weeks of your procedure date, please call us at 410-224-3636.</p>	<p>Stop consuming all food with seeds, corn and nuts 5 days prior to your procedure.</p> <p>Stop taking all NSAIDS. Ibuprofen, Advil, Aleve, etc. 5 days prior to your procedure. You may take Tylenol as needed.</p> <p>Inhalers and Nasal Spray- Continue as prescribed. Please bring with you on day of procedure.</p> <p>Blood Pressure and other essential medications: Continue as prescribed. Take these medications on the day of your procedure with a SMALL sip of water at least 2 hours before procedure.</p> <p>If you use a CPAP: please bring with you on the day of you procedure.</p> <p>If you wear contacts: please remove them at home or bring your contact lens supplies with you if you don't have glasses to wear.</p> <p>Please leave jewelry and other valuables at home.</p> <p>Contact our pre-operative department with any questions at 410 224-3636 Option 2.</p> <p>3 days prior – no solid food after midnight</p>	<p>Clear liquid diet all day today.</p> <p>Acceptable clear liquids include: water, clear broth, apple juice, white cranberry juice, white grape juice, soda, jello, popsicles, coffee, and tea. You may have any transparent liquid. Nothing with milk or creamer.</p> <p>Nothing red, blue or purple in color are allowed.</p> <p>6pm: Clenpiq Prep- (Part 1)</p> <p>Step 1-Drink 1st bottle of Clenpiq solution. Shake well prior to drinking. Drink all of the solution in the bottle.</p> <p>Step 2- Drink five 8oz glasses of clear liquid (total of 40oz). Drink each 8oz every 15 minutes. It should take you no more than 1 hour to complete these steps.</p> <p>After completing this part of the prep take 10 mg of Bisacodyl (2 tablets –5mg each.</p> <p>If you are unable to tolerate the prep, please call us.</p>	<p>Clear liquid diet continues all day today.</p> <p>7am: Clenpiq Prep- (Part 2)</p> <p>Step 1-Drink 1st bottle of Clenpiq solution. Shake well prior to drinking. Drink all of the solution in the bottle.</p> <p>Step 2- Drink five 8oz glasses of clear liquid (total of 40oz). Drink each 8oz every 15 minutes.</p> <p>It should take you no more than 1 hour to complete these steps.</p> <p>6pm: Plenvu Prep- (Part 1) :</p> <p>Step 1-Empty Dose 1 Pouch in container. Add 16oz. water. Mix with spoon to dissolve.</p> <p>Step 2- Drink solution until all solution has been consumed within 30 minutes.</p> <p>Step 3- Refill container with 16oz water and drink slowly over 30 minutes.</p> <p>After completing this part of the prep take 10 mg of Bisacodyl (2 tablets –5mg each.</p> <p>If your prep results are not getting to a transparent liquid, please call us.</p>	<p>Clear liquid diet until 4 hours prior to your procedure.</p> <p>Morning Dose of Clenpiq: (Part 2) 5 hours prior to your procedure.</p> <p>Step 1-Drink 2nd bottle of Clenpiq solution. Shake well prior to drinking. Drink all of the solution in the bottle.</p> <p>Step 2- Drink five 8oz glasses of clear liquid (total of 40oz). Drink each 8oz every 15 minutes. It should take you no more than 1 hour to complete these steps.</p> <p>You must finish all of the prep, including the water 4 hours prior to your procedure. Please call the office if your prep results are not clear like water or a transparent yellow at this time.</p> <p>You cannot have ANYTHING AT ALL by mouth for at least 4 hours prior to scheduled procedure time. This includes ice, water, gum, hard candy and mints.</p> <p>No smoking the day of procedure.</p> <p>Please remember to bring your:</p> <ul style="list-style-type: none"> -Current Medication list with the last dose for each medication taken -Insurance Cards -Driver's License/Photo ID -Any co-insurance fees that are due.



MDTEC Medication Record Form

- This form must be **completed** prior to your procedure.
- Please list all prescriptions, over the counter medications, vitamins and supplements taken in the last month.
- Please remember to review medication instructions provided on page 1 of your procedure instructions.
- *If you have questions regarding your medications prior to your procedure, please call 410-224-3636 and ask to speak with our procedure coordinator.*

Medication Name	Dose	Frequency and Time(s) Taken	<u>Date / Time of Last Dose</u>

Patient Signature

Date

