

Colonoscopy Prep Instructions – Colyte & Bisacodyl & 2 doses of Milk of Magnesia (Colyte also called Gavilyte or Peg 3350 or TriLyte or Nulytley)



YOUR BOWEL PREP IS VERY IMPORTANT - Please read immediately and review instructions carefully! Only follow these MDTEC instructions for your prep.

If you do not follow these instructions and you are not prepped properly for your procedure, a cancellation fee will apply.

Prior to your procedure:	Read 7 Days Prior	2 Days Prior	1 Day Prior	Procedure Day
Please pick up your prescription within 72 hours. AND- Purchase Bisacodyl 5mg tablets (4) & 1 bottle of Milk of Magnesia. You will need to arrange for a responsible adult to drive you home. You cannot drive until the day after your procedure. A taxi, ride share, or shuttle is not an approved means of transportation unless you have a family member or friend with you. If you take Phentermine – stop taking this 10 days prior to your procedure. If you take weekly injections for weight loss or Diabetes: do NOT take this within 7 days of your procedure. If you take iron, dietary, herbal or fiber supplements, stop taking these 7 days prior to your procedure. If you take a blood thinner- Please contact the prescribing doctor to obtain a blood thinner hold order. The office can fax this order to us at 410-972-2698. If you see a specialist: you can request that your most recent visit summary, labs, and test results from that office be faxed to us at 410-972-2698 If you develop any covid, flu, or cold like symptoms within 2 weeks of your procedure date, please call us at 410-224-3636.	Avoid consuming all food with seeds, corn and nuts 5 days prior to your procedure. Stop taking all NSAIDS. Ibuprofen, Advil, Aleve, etc. 5 days prior to your procedure. You may take Tylenol as needed. Inhalers and Nasal Spray- Continue as prescribed. Please bring with you on day of procedure. Blood Pressure and other essential medications: Continue as prescribed. Take these medications on the day of your procedure with a SMALL sip of water at least 2 hours before procedure. If you wear contacts: please remove them at home or bring your contact lens supplies with you if you don't have glasses to wear. Please leave jewelry and other valuables at home. Contact our pre-operative department with any questions at 410 224-3636 Option 2.	Drink at least four 8oz. glasses of water throughout the day.  Eat a light dinner. 6pm: Begin clear liquid diet. Take 4 tablespoons of Milk of Magnesia, followed by 8 ounces of water. Acceptable clear liquids include: water, clear broth, apple juice, white cranberry juice, white grape juice, soda, jello, popsicles, coffee, and tea. You may have any transparent liquid. Nothing with milk or creamer. Nothing red, blue or purple in color are allowed.	Clear liquid diet all day today. Drink at least four 8oz. glasses of water throughout the day.  No alcohol the day before or day of your procedure. If you are taking diabetic medication, cut your dose in half this day and do not take any diabetic medication day of procedure. 4pm: Take 4 tablespoons of Milk of Magnesia, followed by 8 ounces of water. 6pm: Colyte Prep- (Part 1) Step 1 - Add cool water to line on container. Close and shake to dissolve. Add Flavor Pack to Container if not already added. Step 2 - Drink 8oz of solution every 15 minutes until 75% (3/4) has been consumed. You should be done by ~9pm. After completing this part of the prep take 10 mg of Bisacodyl (2 tablets – 5mg each). Step 3 - Store the remainder of the solution in the refrigerator overnight. If you are unable to tolerate the prep, please call 410-224-4887 to speak to the on call doctor.	Clear liquid diet until 4 hours prior to your procedure. Morning Dose Colyte (Part 2) 5 hours prior to your procedure: Drink 8oz of solution every 15 minutes until rest of solution has been consumed. This should be completed within one hour. After completing this part of the prep take 10 mg of Bisacodyl (2 tablets – 5mg each). You must finish all of the prep, including the water 4 hours prior to your procedure. Please call the office if your prep results are not clear like water or a transparent yellow at this time. You cannot have ANYTHING AT ALL by mouth for at least 4 hours prior to scheduled procedure time. This includes ice, water, gum, hard candy and mints. No smoking the day of procedure. Please remember to bring your: -Current Medication list with the last dose for each medication taken -Insurance Cards -Driver's License/Photo ID -Any co-insurance fees that are due.



MDTEC

Medication Record Form

- This form must be completed prior to your procedure.
- Please list all prescriptions, over the counter medications, vitamins and supplements taken in the last month.
- Please remember to review medication instructions provided on page 1 of your procedure instructions.
- *If you have questions regarding your medications prior to your procedure, please call 410-224-3636 and ask to speak with our procedure coordinator.*

Medication Name	Dose	Frequency and Time(s) Taken	<u>Date / Time of Last Dose</u>

Patient Signature

Date



MDTEC

Medication Record Form

Patient Signature

Date