Colonoscopy Prep Instructions – Moviprep with Bisacodyl & 2 doses of Milk of Magnesia



621 Ridgley Ave, Suite 101 Annapolis, MD 21401 410-224-3636

F: 410-972-2698

YOUR BOWEL Prep IS VERY IMPORTANT- Please read immediately and review instructions carefully! Only follow these MDTEC instructions for your prep. If you do not follow these instructions and you are not prepped properly for your procedure, a cancellation fee will apply.

If you do not follow these instructions and you are not prepped properly for your procedure, a cancellation fee will apply.								
Prior to your procedure:	Read 7 Days Prior	2 Days Prior	1 Day Prior	Procedure Day				
Please pick up your prescription	Avoid consuming all food	Drink at least four	Clear liquid diet continues all day	Clear liquid diet until 4 hours prior to				
within 72 hours.	with seeds, corn and nuts	8oz. glasses of	today.	<mark>your procedure.</mark>				
AND-Purchase Bisacodyl 5mg	<u>5 days</u> prior to your	water throughout	Drink at least four 8oz. glasses of	Morning dose of MoviPrep - Part 2				
tablets (4) and 1 bottle of Milk of	proc <mark>edure.</mark>	the day.	water throughout the day.	5 hours prior to your procedure.				
Magnesia.	Stop taking all NSAIDS.	\ <u>\</u>	\(\(\(\(\(\(\(\(\(_	Step 1-Empty Pouches (A&B) into the				
You will need to arrange for a	Ibuprofen, Advil, Aleve,			container. Add warm water to top line <u>on</u>				
responsible adult to drive you home.	etc.	Eat a light dinner.	No alcohol the day before or day of	the container. Mix to dissolve.				
You cannot drive until the day after	5 days prior to your	6pm:	your procedure.	Step 2- Container has 4 marks. Drink 8oz				
your procedure.	procedure. You may take Tylenol as needed.		If you are taking diabetic	of solution (to the mark) every 15 minutes				
A taxi, ride share, or shuttle is not an	Tylenor as needed.	Begin Clear liquid	medication, cut your dose in half	until all solution has been consumed.				
approved means of transportation	Inhalers and Nasal Spray-	diet.	this day and do not take any	It should take you no more than 1 hour				
unless you have a family member or	Continue as prescribed.	Take 4	diabetic medication day of	to complete these steps.				
friend with you.	Please bring with you on	tablespoons of	procedure.	After completing this part of the prep take				
If you take Phentermine: stop taking	day of procedure.	Milk of Magnesia	4pm: Take 4 tablespoons of Milk	10 mg of Bisacodyl (2 tablets –5mg each)				
this 10 days prior to your procedure.	Blood Pressure and other	followed by 8oz	of Magnesia followed by 8oz of					
If you take weekly injections for	essential medications:	of water	water	You must finish all of the prep, including				
weight loss or Diabetes: do NOT take	Continue as prescribed.	Acceptable clear		the water 4 hours prior to your				
this within 7 days of your procedure.	Take these medications	liquids include:	6pm: MoviPrep - Part 1	procedure. Please call the office if your prep results are not clear like water or a				
If you take iron, dietary, herbal or	on the day of your	water, clear broth,	Step 1-Empty Pouches (A&B) into	transparent yellow at this time.				
fiber supplements: stop taking these 7	procedure with a SMALL	apple juice, white	the container. Add warm water to	transparent yenow at tins time.				
days prior to your procedure.	sip of water at least 2	cranberry juice,	top line on the container. Mix to	You cannot have ANYTHING AT ALL by				
If you take a blood thinner:	hours before procedure.	white grape juice,	dissolve.	mouth for at least 4 hours prior to				
Please contact the prescribing doctor	If you wear contacts:	soda, jello,	Step 2 - Container has 4 marks.	scheduled procedure time. This includes				
to obtain a blood thinner hold order.	please remove them at	popsicles, coffee,	Drink 8oz of solution (to the mark)	ice, water, gum, hard candy and mints.				
The office can fax this order to us at	home or bring your	and tea.	every 15 minutes until all solution					
410-972-2698.	contact lens supplies with	You may have any	has been consumed.	No smoking the day of procedure.				
If you see a specialist: We will need to	you if you don't have	transparent liquid.	After completing this part of the prep					
review the notes from your most	glasses to wear.	Nothing with milk	take 10 mg of Bisacodyl (2 tablets -	Please remember to bring your:				
recent office visit, labs, and test results prior to your procedure. You can call	Please leave jewelry and	or creamer.	5mg each)	-Current Medication list with the last				
their office to request these be faxed	other valuables at home.	Nothing red, blue	It should take you no more than 1	dose for each medication taken				
to us at 410-972-2698.		or purple in color	hour to complete these steps.	-Insurance Cards -Driver's License/Photo ID				
If you develop any covid, flu, or cold	Contact our pre-	are allowed.	If you are unable to tolerate the	-Driver's License/Photo ID -Any co-insurance fees that are due.				
like symptoms within 2 weeks of your	operative department		prep, please call 410-224-4887 to	-Any co-mountaince nees that are due.				
procedure date, please call us at 410-	with any questions at		speak to the on call doctor.					
224-3636	410 224-3636 Option 2		-					



- This form must be <u>completed</u> prior to your procedure.
- Please list all prescriptions, over the counter medications, vitamins and supplements taken in the last month.
- Please remember to review medication instructions provided on page 1 of your procedure instructions.
- If you have questions regarding your medications prior to your procedure, please call 410-224-3636 and ask to speak with our procedure coordinator.

Medication Name	Dose	Frequency and Time(s) Taken	Date / Time of Last Dose

Patient Signature	Date



Patient Signature

Date