



Colonoscopy Prep Instructions – Plenvu with 1 Bottle of Magnesium Citrate



MARYLAND DIAGNOSTIC & THERAPEUTIC ENDO CENTER
 621 Ridgley Ave, Suite 101
 Annapolis, MD 21401
 410-224-3636
 F: 410-972-2698

YOUR BOWEL PREP IS VERY IMPORTANT - Please read immediately and review instructions carefully! Only follow these MDTEC instructions for your prep.

If you do not follow these instructions and you are not prepped properly for your procedure, a cancellation fee will apply.

Prior to your procedure:		2 Days Prior	1 Day Prior	Procedure Day
<p>Your prescription is sent to your pharmacy after your pre-op call or office visit has been completed. Please pick up your prescription within 72 hours.</p> <p>AND-Purchase 1 bottle of Magnesium Citrate.</p> <p>You will need to arrange for a responsible adult to drive you home. You cannot drive until the day after your procedure. A taxi, ride share, or shuttle is not an approved means of transportation unless you have a family member or friend with you.</p> <p>If you take Phentermine: stop taking this 10 days prior to your procedure.</p> <p>If you take weekly injections for weight loss or Diabetes: do NOT take this within 7 days of your procedure.</p> <p>If you take iron, dietary, herbal or fiber supplements: stop taking these 7 days prior to your procedure.</p> <p>If you take a blood thinner:</p> <p>Please contact the prescribing doctor to obtain a blood thinner hold order. The office can fax this order to us at 410-972-2698.</p> <p>If you see a specialist: We will need to review the notes from your most recent office visit, labs, and test results prior to your procedure. You can call their office to request these be faxed to us at 410-972-2698.</p> <p>If you develop any covid, flu, or cold like symptoms within 2 weeks of your procedure date, please call us at 410-224-3636</p>	<p>Avoid consuming all food with seeds, corn and nuts 5 days prior to your procedure.</p> <p>Stop taking all NSAIDS. Ibuprofen, Advil, Aleve, etc. 5 days prior to your procedure. You may take Tylenol as needed.</p> <p>Inhalers and Nasal Spray- Continue as prescribed. Please bring with you on day of procedure.</p> <p>Blood Pressure and other essential medications: Continue as prescribed. Take these medications on the day of your procedure with a SMALL sip of water at least 2 hours before procedure.</p> <p>If you wear contacts: please remove them at home or bring your contact lens supplies with you if you don't have glasses to wear. Please leave jewelry and other valuables at home.</p> <p>Contact our pre-operative department with any questions at 410 224-3636 Option 2.</p>	<p>Drink at least four 8oz. glasses of water throughout the day.</p>  <p>Eat a light dinner.</p> <p>6pm: Begin clear liquid diet.</p> <p>Drink 1st Bottle of Magnesium Citrate. Then Drink One 8oz glass of Water.</p> <p>Acceptable clear liquids include: water, clear broth, apple juice, white cranberry juice, white grape juice, soda, jello, popsicles, coffee, and tea. You may have any transparent liquid. Nothing with milk or creamer.</p> <p>Nothing red, blue or purple in color are allowed.</p>	<p>Clear liquid diet continues all day today.</p> <p>Drink at least four 8oz. glasses of water throughout the day.</p>  <p>No alcohol the day before or day of your procedure.</p> <p>If you are taking diabetic medication, cut your dose in half this day and do not take any diabetic medication day of procedure.</p> <p>6pm: Plenvu Prep- (Part 1) Step 1-Empty Dose 1 Pouch in container. Add 16oz. water. Mix with spoon to dissolve. Step 2- Drink solution until all solution has been consumed within 30 minutes. Step 3- Refill container with 16oz water and drink slowly over 30 minutes.</p> <p>It should take you no more than 1 hour to complete these steps.</p> <p>If you are unable to tolerate the prep, please call 410-224-4887 to speak to the on call doctor.</p>	<p>Clear liquid diet until 4 hours prior to your procedure.</p> <p>Morning Dose of Plenvu: (Part 2) 5 hours prior to your procedure. Step 1- Empty Dose 2 Pouch A & B in container. Add 16oz. water. Mix with spoon to dissolve. Step 2- Drink solution until all solution has been consumed within 30 minutes. Step 3- Refill container with 16oz water and drink slowly over 30 minutes.</p> <p>You must finish all of the prep, including the water 4 hours prior to your procedure. Please call the office if your prep results are not clear like water or a transparent yellow at this time.</p> <p>You cannot have ANYTHING AT ALL by mouth for at least 4 hours prior to scheduled procedure time. This includes ice, water, gum, hard candy and mints.</p> <p>No smoking the day of procedure. Please remember to bring your:</p> <ul style="list-style-type: none"> -Current Medication list with the last dose for each medication taken -Insurance Cards -Driver's License/Photo ID -Any co-insurance fees that are due.

